

# Recall: $\lambda$ -Terms, $M$

are built up from a given, countable collection of

- ▶ variables  $x, y, z, \dots$

by two operations for forming  $\lambda$ -terms:

- ▶  $\lambda$ -abstraction:  $(\lambda x.M)$   
(where  $x$  is a variable and  $M$  is a  $\lambda$ -term)
- ▶ application:  $(M M')$   
(where  $M$  and  $M'$  are  $\lambda$ -terms).

# $\beta$ -Reduction

Recall that  $\lambda x.M$  is intended to represent the function  $f$  such that  $f(x) = M$  for all  $x$ . We can regard  $\lambda x.M$  as a function on  $\lambda$ -terms via substitution: map each  $N$  to  $M[N/x]$ .

↑ result of substituting  
 $N$  for free  $x$  in  $M$

# Substitution $N[M/x]$

$$\begin{aligned}x[M/x] &= M \\y[M/x] &= y \quad \text{if } y \neq x \\(\lambda y.N)[M/x] &= \lambda y.N[M/x] \quad \text{if } y \# (M x) \\(N_1 N_2)[M/x] &= N_1[M/x] N_2[M/x]\end{aligned}$$

$N[M/x]$  = result of replacing all free occurrences of  $x$  in  $N$  with  $M$ , avoiding "capture" of free variables in  $M$  by  $\lambda$ -binders in  $N$

# Substitution $N[M/x]$

$$\begin{aligned}x[M/x] &= M \\y[M/x] &= y \quad \text{if } y \neq x \\(\lambda y.N)[M/x] &= \lambda y.N[M/x] \quad \text{if } y \# (M x) \\(N_1 N_2)[M/x] &= N_1[M/x] N_2[M/x]\end{aligned}$$

Side-condition  $y \# (M x)$  ( $y$  does not occur in  $M$  and  $y \neq x$ ) makes substitution “capture-avoiding”.

E.g. if  $x \neq y$

$$(\lambda y.x)[y/x] \neq \lambda y.y$$

# Substitution $N[M/x]$

$$\begin{aligned}x[M/x] &= M \\y[M/x] &= y \quad \text{if } y \neq x \\(\lambda y.N)[M/x] &= \lambda y.N[M/x] \quad \text{if } y \# (M x) \\(N_1 N_2)[M/x] &= N_1[M/x] N_2[M/x]\end{aligned}$$

Can always satisfy this  up to  $\alpha$ -equivalence

E.g. if  $x \neq y \neq z \neq x$

$$(\lambda y.x)[y/x] =_{\alpha} (\lambda z.x)[y/x] = \lambda z.y$$

In fact  $N \mapsto N[M/x]$  induces a totally defined function from the set of  $\alpha$ -equivalence classes of  $\lambda$ -terms to itself.

$$\lambda x. (\lambda z. z) y x \quad [ \lambda z. y / y ]$$

||

$$\lambda x. (\lambda z. z) y x \quad [ \lambda z. y / y ]$$

no possible  
capture

||

$$\begin{aligned} & \lambda x. (\lambda z. z) y x \left[ \lambda z. y / y \right] \\ &= \lambda x. (\lambda z. z) (\lambda z. y) x \end{aligned}$$

---

$$\begin{aligned} & \lambda x. (\lambda u. u) x y \left[ \lambda y. x / y \right] \\ &= \end{aligned}$$

$$\lambda x. (\lambda z. z) y x \left[ \lambda x. y / y \right]$$
$$= \lambda x. (\lambda z. z) (\lambda x. y) x$$

---

$$\lambda x. (\lambda u. u) x y \left[ \lambda y. x / y \right] \quad \text{possible capture}$$
$$=$$

$$\lambda x. (\lambda z. z) y x \ [ \lambda x. y / y ]$$

$$= \lambda x. (\lambda z. z) (\lambda x. y) x$$


---

$$\lambda x. (\lambda u. u) x y \ [ \lambda y. x / y ]$$

possible capture...

$$=_{\alpha} \lambda z. (\lambda u. u) z y \ [ \lambda y. x / y ]$$

... $\alpha$ -convert to avoid

$$\lambda x. (\lambda z. z) y x \left[ \lambda x. y / y \right]$$

$$= \lambda x. (\lambda z. z) (\lambda x. y) x$$


---

$$\lambda x. (\lambda u. u) x y \left[ \lambda y. x / y \right] \quad \text{possible capture...}$$

$$\stackrel{\alpha}{=} \lambda z. (\lambda u. u) z y \left[ \lambda y. x / y \right] \quad \text{...}\alpha\text{-convert to avoid}$$

$$= \lambda z. (\lambda u. u) z (\lambda y. x)$$

# $\beta$ -Reduction

Recall that  $\lambda x.M$  is intended to represent the function  $f$  such that  $f(x) = M$  for all  $x$ . We can regard  $\lambda x.M$  as a function on  $\lambda$ -terms via substitution: map each  $N$  to  $M[N/x]$ .

So the natural notion of computation for  $\lambda$ -terms is given by stepping from a

$\beta$ -redex  $(\lambda x.M)N$

to the corresponding

$\beta$ -reduct  $M[N/x]$


# $\beta$ -Reduction

One-step  $\beta$ -reduction,  $M \rightarrow M'$ :

$$\frac{}{(\lambda x.M)N \rightarrow M[N/x]} \qquad \frac{M \rightarrow M'}{\lambda x.M \rightarrow \lambda x.M'}$$
$$\frac{M \rightarrow M'}{MN \rightarrow M'N} \qquad \frac{M \rightarrow M'}{NM \rightarrow NM'}$$
$$\frac{N =_{\alpha} M \quad M \rightarrow M' \quad M' =_{\alpha} N'}{N \rightarrow N'}$$


# $\beta$ -Reduction

E.g.




# $\beta$ -Reduction

E.g.




# $\beta$ -Reduction

E.g.




# $\beta$ -Reduction

E.g.



# $\beta$ -Reduction

E.g.



Many-step  $\beta$ -reduction,  $M \twoheadrightarrow M'$ :

$$\frac{M =_{\alpha} M'}{M \twoheadrightarrow M'} \quad \left[ \frac{M \rightarrow M'}{M \twoheadrightarrow M'} \right] \quad \frac{M \twoheadrightarrow M' \quad M' \rightarrow M''}{M \twoheadrightarrow M''}$$

(no steps)                      (1 step)                      (1 more step)

E.g.

$$(\lambda x.x y)((\lambda y z.z)u) \twoheadrightarrow y$$

# $\beta$ -Conversion $M =_{\beta} N$

Informally:  $M =_{\beta} N$  holds if  $N$  can be obtained from  $M$  by performing zero or more steps of  $\alpha$ -equivalence,  $\beta$ -reduction, or  $\beta$ -expansion (= inverse of a reduction).

E.g.  $u((\lambda x y. v x)y) =_{\beta} (\lambda x. u x)(\lambda x. v y)$

because  $(\lambda x. u x)(\lambda x. v y) \rightarrow u(\lambda x. v y)$

and so we have

$u((\lambda x y. v x)y) =_{\alpha} u((\lambda x y'. v x)y)$

# $\beta$ -Conversion $M =_{\beta} N$

Informally:  $M =_{\beta} N$  holds if  $N$  can be obtained from  $M$  by performing zero or more steps of  $\alpha$ -equivalence,  $\beta$ -reduction, or  $\beta$ -expansion (= inverse of a reduction).

E.g.  $u((\lambda x y. v x)y) =_{\beta} (\lambda x. u x)(\lambda x. v y)$

because  $(\lambda x. u x)(\lambda x. v y) \rightarrow u(\lambda x. v y)$

and so we have

$$\begin{array}{lcl} u((\lambda x y. v x)y) & =_{\alpha} & u((\lambda x y'. v x)y) \\ & \rightarrow & u(\lambda y'. v y) \end{array} \quad \text{reduction}$$

# $\beta$ -Conversion $M =_{\beta} N$

Informally:  $M =_{\beta} N$  holds if  $N$  can be obtained from  $M$  by performing zero or more steps of  $\alpha$ -equivalence,  $\beta$ -reduction, or  $\beta$ -expansion (= inverse of a reduction).

E.g.  $u((\lambda x y. v x)y) =_{\beta} (\lambda x. u x)(\lambda x. v y)$

because  $(\lambda x. u x)(\lambda x. v y) \rightarrow u(\lambda x. v y)$

and so we have

$$\begin{aligned} u((\lambda x y. v x)y) &=_{\alpha} u((\lambda x y'. v x)y) \\ &\rightarrow u(\lambda y'. v y) && \text{reduction} \\ &=_{\alpha} u(\lambda x. v y) \end{aligned}$$

# $\beta$ -Conversion $M =_{\beta} N$

Informally:  $M =_{\beta} N$  holds if  $N$  can be obtained from  $M$  by performing zero or more steps of  $\alpha$ -equivalence,  $\beta$ -reduction, or  $\beta$ -expansion (= inverse of a reduction).

E.g.  $u((\lambda x y. v x)y) =_{\beta} (\lambda x. u x)(\lambda x. v y)$

because  $(\lambda x. u x)(\lambda x. v y) \rightarrow u(\lambda x. v y)$

and so we have

$$\begin{aligned} u((\lambda x y. v x)y) &=_{\alpha} u((\lambda x y'. v x)y) \\ &\rightarrow u(\lambda y'. v y) && \text{reduction} \\ &=_{\alpha} u(\lambda x. v y) \\ &\leftarrow (\lambda x. u x)(\lambda x. v y) && \text{expansion} \end{aligned}$$

# $\beta$ -Conversion $M =_{\beta} N$

is the binary relation inductively generated by the rules:

$$\frac{M =_{\alpha} M'}{M =_{\beta} M'}$$

$$\frac{M \rightarrow M'}{M =_{\beta} M'}$$

$$\frac{M =_{\beta} M'}{M' =_{\beta} M}$$

$$\frac{M =_{\beta} M' \quad M' =_{\beta} M''}{M =_{\beta} M''}$$

$$\frac{M =_{\beta} M'}{\lambda x.M =_{\beta} \lambda x.M'}$$

$$\frac{M =_{\beta} M' \quad N =_{\beta} N'}{MN =_{\beta} M'N'}$$

# Church-Rosser Theorem

**Theorem.**  $\rightarrow$  is **confluent**, that is, if  $M_1 \leftarrow M \rightarrow M_2$ , then there exists  $M'$  such that  $M_1 \rightarrow M' \leftarrow M_2$ .

[Proof omitted.]

# Church-Rosser Theorem

**Theorem.**  $\rightarrow$  is **confluent**, that is, if  $M_1 \leftarrow M \rightarrow M_2$ , then there exists  $M'$  such that  $M_1 \rightarrow M' \leftarrow M_2$ .

**Corollary.** To show that two terms are  $\beta$ -convertible, it suffices to show that they both reduce to the same term. More precisely:  $M_1 =_{\beta} M_2$  iff  $\exists M (M_1 \rightarrow M \leftarrow M_2)$ .

# Church-Rosser Theorem

**Theorem.**  $\rightarrow$  is **confluent**, that is, if  $M_1 \leftarrow M \rightarrow M_2$ , then there exists  $M'$  such that  $M_1 \rightarrow M' \leftarrow M_2$ .

**Corollary.**  $M_1 =_{\beta} M_2$  iff  $\exists M (M_1 \rightarrow M \leftarrow M_2)$ .

**Proof.**  $=_{\beta}$  satisfies the rules generating  $\rightarrow$ ; so  $M \rightarrow M'$  implies  $M =_{\beta} M'$ . Thus if  $M_1 \rightarrow M \leftarrow M_2$ , then  $M_1 =_{\beta} M =_{\beta} M_2$  and so  $M_1 =_{\beta} M_2$ .

Conversely,

# Church-Rosser Theorem

**Theorem.**  $\rightarrow\rangle$  is **confluent**, that is, if  $M_1 \leftarrow M \rightarrow\rangle M_2$ , then there exists  $M'$  such that  $M_1 \rightarrow\rangle M' \leftarrow M_2$ .

**Corollary.**  $M_1 =_\beta M_2$  iff  $\exists M (M_1 \rightarrow\rangle M \leftarrow M_2)$ .

**Proof.**  $=_\beta$  satisfies the rules generating  $\rightarrow\rangle$ ; so  $M \rightarrow\rangle M'$  implies  $M =_\beta M'$ . Thus if  $M_1 \rightarrow\rangle M \leftarrow M_2$ , then  $M_1 =_\beta M =_\beta M_2$  and so  $M_1 =_\beta M_2$ .

Conversely, the relation  $\{(M_1, M_2) \mid \exists M (M_1 \rightarrow\rangle M \leftarrow M_2)\}$  satisfies the rules generating  $=_\beta$ : the only difficult case is closure of the relation under transitivity and for this we use the Church-Rosser theorem:  $M_1 \rightarrow\rangle M \leftarrow M_2 \rightarrow\rangle M' \leftarrow M_3$



# Church-Rosser Theorem

**Theorem.**  $\rightarrow\rightarrow$  is **confluent**, that is, if  $M_1 \leftarrow M \rightarrow\rightarrow M_2$ , then there exists  $M'$  such that  $M_1 \rightarrow\rightarrow M' \leftarrow M_2$ .

**Corollary.**  $M_1 =_{\beta} M_2$  iff  $\exists M (M_1 \rightarrow\rightarrow M \leftarrow M_2)$ .

**Proof.**  $=_{\beta}$  satisfies the rules generating  $\rightarrow\rightarrow$ ; so  $M \rightarrow\rightarrow M'$  implies  $M =_{\beta} M'$ . Thus if  $M_1 \rightarrow\rightarrow M \leftarrow M_2$ , then  $M_1 =_{\beta} M =_{\beta} M_2$  and so  $M_1 =_{\beta} M_2$ .

Conversely, the relation  $\{(M_1, M_2) \mid \exists M (M_1 \rightarrow\rightarrow M \leftarrow M_2)\}$  satisfies the rules generating  $=_{\beta}$ : the only difficult case is closure of the relation under transitivity and for this we use the Church-Rosser theorem. Hence  $M_1 =_{\beta} M_2$  implies  $\exists M (M_1 \rightarrow\rightarrow M' \leftarrow M_2)$ .

# $\beta$ -Normal Forms

**Definition.** A  $\lambda$ -term  $N$  is in  $\beta$ -normal form (nf) if it contains no  $\beta$ -redexes (no sub-terms of the form  $(\lambda x.M)M'$ ).  $M$  has  $\beta$ -nf  $N$  if  $M =_{\beta} N$  with  $N$  a  $\beta$ -nf.

# $\beta$ -Normal Forms

**Definition.** A  $\lambda$ -term  $N$  is in  $\beta$ -normal form (nf) if it contains no  $\beta$ -redexes (no sub-terms of the form  $(\lambda x.M)M'$ ).  $M$  has  $\beta$ -nf  $N$  if  $M =_{\beta} N$  with  $N$  a  $\beta$ -nf.

Note that if  $N$  is a  $\beta$ -nf and  $N \rightarrow N'$ , then it must be that  $N =_{\alpha} N'$  (why?).

Hence if  $N_1 =_{\beta} N_2$  with  $N_1$  and  $N_2$  both  $\beta$ -nfs, then  $N_1 =_{\alpha} N_2$ . (For if  $N_1 =_{\beta} N_2$ , then by Church-Rosser  $N_1 \rightarrow M' \leftarrow N_2$  for some  $M'$ , so  $N_1 =_{\alpha} M' =_{\alpha} N_2$ .)

**So the  $\beta$ -nf of  $M$  is unique up to  $\alpha$ -equivalence if it exists.**

(and if  $M$  does have  $\beta$ -nf  $N$ , then  
 $M \rightarrow N$ )

# Non-termination

Some  $\lambda$  terms have no  $\beta$ -nf.

E.g.  $\Omega \triangleq (\lambda x.x x)(\lambda x.x x)$  satisfies

- ▶  $\Omega \rightarrow (x x)[(\lambda x.x x)/x] = \Omega$ ,
- ▶  $\Omega \rightarrow M$  implies  $\Omega =_{\alpha} M$ .

So there is no  $\beta$ -nf  $N$  such that  $\Omega =_{\beta} N$ .

# Non-termination

Some  $\lambda$  terms have no  $\beta$ -nf.

E.g.  $\Omega \triangleq (\lambda x.x x)(\lambda x.x x)$  satisfies

- ▶  $\Omega \rightarrow (x x)[(\lambda x.x x)/x] = \Omega$ ,
- ▶  $\Omega \twoheadrightarrow M$  implies  $\Omega =_{\alpha} M$ .

So there is no  $\beta$ -nf  $N$  such that  $\Omega =_{\beta} N$ .

**A term can possess both a  $\beta$ -nf and infinite chains of reduction from it.**

E.g.  $(\lambda x.y)\Omega \rightarrow y$ , but also  $(\lambda x.y)\Omega \rightarrow (\lambda x.y)\Omega \rightarrow \dots$ .

# Non-termination

**Normal-order reduction** is a deterministic strategy for reducing  $\lambda$ -terms: reduce the “left-most, outer-most” redex first.

- ▶ left-most: reduce  $M$  before  $N$  in  $M N$ , and then
- ▶ outer-most: reduce  $(\lambda x.M)N$  rather than either of  $M$  or  $N$ .

(cf. call-by-name evaluation).

**Fact:** normal-order reduction of  $M$  always reaches the  $\beta$ -nf of  $M$  if it possesses one.

$$\frac{M_1 =_{\alpha} M'_1 \quad M'_1 \rightarrow_{\text{NOR}} M'_2 \quad M'_2 =_{\alpha} M_2}{M_1 \rightarrow_{\text{NOR}} M_2}$$

$$\frac{M \rightarrow_{\text{NOR}} M'}{\lambda x. M \rightarrow_{\text{NOR}} \lambda x. M'}$$

$$\frac{M_1 \rightarrow_{\text{NOR}} M'_1}{M_1 M_2 \rightarrow_{\text{NOR}} M'_1 M_2}$$

$$\frac{}{(\lambda x. M) M' \rightarrow_{\text{NOR}} M[M'/x]}$$

$$\frac{M \rightarrow_{\text{NOR}} M'}{UM \rightarrow_{\text{NOR}} UM'}$$

where

$$\begin{cases} U ::= x \mid UN \\ N ::= \lambda x. N \mid U \end{cases}$$

$\beta$ -normal forms

"neutral" forms