

Topics in Concurrency

Lectures 5

Glynn Winskel

18 February 2019

Specification logics

Logics for specifying correctness properties.

We'll look at:

- Basic logics and bisimilarity
- Fixed points and logic
- CTL
- Model checking

Finitary Hennessy-Milner Logic

Assertions:

$$A ::= T \mid F \mid A_0 \wedge A_1 \mid A_0 \vee A_1 \mid \neg A \mid \langle \lambda \rangle A \mid \langle - \rangle A \mid [\lambda] A \mid [-] A$$

Satisfaction: $s \models A$

$s \models T$ always

$s \models F$ never

$s \models A_0 \wedge A_1$ if $s \models A_0$ and $s \models A_1$

$s \models A_0 \vee A_1$ if $s \models A_0$ or $s \models A_1$

$s \models \neg A$ if not $s \models A$

$s \models \langle \lambda \rangle A$ if there exists s' s.t. $s \xrightarrow{\lambda} s'$ and $s' \models A$

$s \models \langle - \rangle A$ if there exist s', λ s.t. $s \xrightarrow{\lambda} s'$ and $s' \models A$


$s \models [\lambda] A$ iff for all s' s.t. $s \xrightarrow{\lambda} s'$ have $s' \models A$

$s \models [-] A$ iff for all s', λ s.t. $s \xrightarrow{\lambda} s'$ have $s' \models A$

Alternatively, derived assertions

$$[\lambda] A \equiv \neg \langle \lambda \rangle \neg A \quad [-] A \equiv \neg \langle - \rangle \neg A$$

Examples



? $s \models \langle a \rangle T$?

? $s \models [a] T$?

? $u \models [-] F$?

? $s \models \langle a \rangle \langle b \rangle T$?

? $s \models [a] \langle b \rangle T$?

Examples

Generally:

- $\langle a \rangle T$
- $[a] F$
- $\langle - \rangle F$
- $\langle - \rangle T$
- $[-] T$
- $[-] F$

(Strong) bisimilarity and logic

A non-finitary Hennessy-Milner logic allows an infinite conjunction

$$A ::= \bigwedge_{i \in I} A_i \mid \neg A \mid \langle \lambda \rangle A$$

with semantics

$$s \models \bigwedge_{i \in I} A_i \text{ iff } s \models A_i \text{ for all } i \in I$$

Define

$$p \simeq q \text{ iff } \text{for all assertions } A \text{ of H-M logic} \\ p \models A \text{ iff } q \models A$$

Theorem

$$\simeq = \sim$$

This gives a way to demonstrate non-bisimilarity of states

Fixed points and model checking

- The finitary H-M logic doesn't allow properties such as
the process never deadlocks
- We can add particular extensions (such as always, never) to the logic (CTL)
- Alternatively, what about defining sets of states 'recursively'? The set of states X that can always do some action satisfies:

$$X = \langle - \rangle T \wedge [-] X$$

- A fixed point equation: $X = \varphi(X)$
- But such equations can have many solutions...

Fixed point equations

- In general, an equation of the form $X = \varphi(X)$ can have many solutions for X .
- Fixed points are important: they represent steady or consistent states
- Range of different fixed point theorems applicable in different contexts e.g.


Theorem (1-dimensional Brouwer's fixed point theorem)

Any continuous function $f : [0, 1] \rightarrow [0, 1]$ has at least one fixed point

(used e.g. in proof of existence of Nash equilibria)

- We'll be interested in fixed points of functions on the powerset lattice \rightsquigarrow Knaster-Tarski fixed point theorem and least and greatest fixed points

Least and greatest fixed points on transition systems: examples



In the above transition system, what are the least and greatest subsets of states X , Y and Z that satisfy:

$$X = X$$

$$Y = \langle - \rangle T \wedge [-] Y$$

$$Z = \neg Z$$

The powerset lattice

- Given a set \mathcal{S} , its powerset is

$$\mathcal{P}(\mathcal{S}) = \{S \mid S \subseteq \mathcal{S}\}$$

- Taking the order on its elements to be inclusion, \subseteq , this forms a complete lattice

We are interested in fixed points of functions of the form

$$\varphi : \mathcal{P}(\mathcal{S}) \rightarrow \mathcal{P}(\mathcal{S})$$

- φ is **monotonic** if $S \subseteq S'$ implies $\varphi(S) \subseteq \varphi(S')$
- a **prefixed point** of φ is a set X satisfying $\varphi(X) \subseteq X$
- a **postfixed point** of φ is a set X satisfying $X \subseteq \varphi(X)$

Knaster-Tarski fixed point theorem for minimum fixed points

Theorem

For monotonic $\varphi : \mathcal{P}(\mathcal{S}) \rightarrow \mathcal{P}(\mathcal{S})$, define

$$m = \bigcap \{X \subseteq \mathcal{S} \mid \varphi(X) \subseteq X\}.$$

Then m is a fixed point of φ and, furthermore, is the least prefixed point:

- 1 $m = \varphi(m)$
- 2 $\varphi(X) \subseteq X$ implies $m \subseteq X$

m is conventionally written

$$\mu X. \varphi(X)$$

Used for inductive definitions: syntax, operational semantics, rule-based programs, model checking

Knaster-Tarski fixed point theorem for maximum fixed points

Theorem

For monotonic $\varphi : \mathcal{P}(S) \rightarrow \mathcal{P}(S)$, define

$$M = \bigcup \{X \subseteq S \mid X \subseteq \varphi(X)\}.$$

Then M is a fixed point of φ and, furthermore, is the greatest postfixed point.

- 1 $M = \varphi(M)$
- 2 $X \subseteq \varphi(X)$ implies $X \subseteq M$

M is conventionally written

$$\nu X. \varphi(X)$$

Used for co-inductive definitions, bisimulation, model checking

(Strong) bisimilarity as a maximum fixed point [§5.2 p68]

Bisimilarity can be viewed as a fixed point \rightsquigarrow model checking algorithms.

Given a relation R (on CCS processes or states of transition systems) define:

$$p \varphi(R) q$$

iff

- 1 $\forall \alpha, p'. \quad p \xrightarrow{\alpha} p' \implies \exists q'. \quad q \xrightarrow{\alpha} q' \ \& \ p' R q'$
- 2 $\forall \alpha, q'. \quad q \xrightarrow{\alpha} q' \implies \exists p'. \quad p \xrightarrow{\alpha} p' \ \& \ p' R q'$

Lemma

$R \subseteq \varphi(R)$ iff R is a (strong) bisimulation.

Hence, by Knaster-Tarski fixed point theorem for maximum fixed points:

Theorem

Bisimilarity is the greatest fixed point of φ .

Theorem

Bisimilarity is the greatest fixed point of φ .

Proof.

$$\sim = \bigcup \{R \mid R \text{ is a bisimulation}\} \quad (1)$$

$$= \bigcup \{R \mid R \subseteq \varphi(R)\} \quad (2)$$

$$= \nu X. \varphi(X) \quad (3)$$

(1) is by definition of \sim

(2) is by Lemma

(3) is by Knaster-Tarski for maximum fixed points: note that φ is monotonic



Question: How is this different from the least fixed point of φ ?

The modal μ -calculus [§4.2 p48]

$$A ::= T \mid F \mid A_0 \wedge A_1 \mid A_0 \vee A_1 \mid \neg A \mid \langle \lambda \rangle A \mid \langle - \rangle A \mid X \mid \nu X.A$$

To guarantee monotonicity (and therefore the existence of the fixed point), require the variable X to occur only **positively** in A in $\nu X.A$. That is, X occurs only under an even number of \neg s.

$$\begin{aligned} s \models \nu X.A & \text{ iff } s \in \nu X.A \\ & \text{ i.e. } s \in \bigcup \{ S \subseteq \mathcal{P} \mid S \subseteq A[S/X] \} \\ & \text{ the maximum fixed point of the monotonic} \\ & \text{ function } S \mapsto A[S/X] \end{aligned}$$

As before, we take

$$[\lambda]A \equiv \neg \langle \lambda \rangle \neg A \quad [-]A \equiv \neg \langle - \rangle \neg A$$

Now also take

$$\mu X.A \equiv \neg \nu X.(\neg A[\neg X/X])$$

Example

Consider the process

$$P \stackrel{\text{def}}{=} a.(a.P + b.c.\mathbf{nil})$$

Which states satisfy

- $\mu X.\langle a \rangle X$
- $\nu X.\langle a \rangle X$
- $\mu X.[a] X$
- $\nu X[a] X$

Approximants

Let $\varphi : \mathcal{P}(\mathcal{S}) \rightarrow \mathcal{P}(\mathcal{S})$ be monotonic.

φ is \cap -continuous iff for all decreasing chains $X_0 \supseteq X_1 \supseteq \dots \supseteq X_n \supseteq \dots$

$$\bigcap_{n \in \omega} \varphi(X_n) = \varphi\left(\bigcap_{n \in \omega} X_n\right)$$

If the set of states \mathcal{S} is finite, continuity certainly holds

Theorem

If $\varphi : \mathcal{P}(\mathcal{S}) \rightarrow \mathcal{P}(\mathcal{S})$ is \cap -continuous:

$$\nu X. \varphi(X) = \bigcap_{n \in \omega} \varphi^n(\mathcal{S})$$

Approximants

Let $\varphi : \mathcal{P}(\mathcal{S}) \rightarrow \mathcal{P}(\mathcal{S})$ be monotonic.

φ is \cup -continuous iff for all increasing chains $X_0 \subseteq X_1 \subseteq \dots \subseteq X_n \subseteq \dots$

$$\bigcup_{n \in \omega} \varphi(X_n) = \varphi\left(\bigcup_{n \in \omega} X_n\right)$$

If the set of states \mathcal{S} is finite, continuity certainly holds

Theorem

If $\varphi : \mathcal{P}(\mathcal{S}) \rightarrow \mathcal{P}(\mathcal{S})$ is \cup -continuous:

$$\mu X. \varphi(X) = \bigcup_{n \in \omega} \varphi^n(\emptyset)$$

Proving interpretations

Proposition

$s \models \mu X. \langle a \rangle T \vee \langle - \rangle X$ in any transition system iff there exists a sequence of transitions from s to a state t where an a -action can occur.

Proving interpretations

Proposition

$s \models \nu X. \langle a \rangle X$ in a finite-state transition system iff there exists an infinite sequence of a -transitions from s .

There are **infinite-state** transition systems where $\varphi(X) = \langle a \rangle X$ is not \sqcap -continuous.

Bisimilarity and modal μ

For finite-state processes, modal- μ can be encoded in infinitary H-M logic

if finite-state processes p and q are bisimilar then they satisfy the same modal- μ assertions

Note that logical equivalence in modal- μ does not generally imply bisimilarity (due to the lack of infinitary conjunction)