

.

Lecture 12

Adjoint functors


The concepts of “category”, “functor” and “natural transformation” were invented by Eilenberg and MacLane in order to formalise “adjoint situations”.

They appear everywhere in mathematics, logic and (hence) computer science.

Examples of adjoint situations that we have already seen...


Given

Set $\rightarrow \Sigma$




monoid
(List Σ , @, nil)


Given



Given




Given



Set

function f



monoid (M, \cdot, e)

there exists a unique monoid morphism with $\hat{f} \circ \eta_\Sigma = f$

$$\hat{f}[a_1, \dots, a_n] = (fa_1) \cdot (fa_2) \cdot \dots \cdot (fa_n)$$

Free monoids

$$\frac{\Sigma \rightarrow U(M, \cdot, e) \text{ morphisms in Set}}{F \Sigma \rightarrow (M, \cdot, e) \text{ morphisms in Mon}}$$

bijection

$$\text{Set}(\Sigma, U(M, \cdot, e)) \cong \text{Mon}(F \Sigma, (M, \cdot, e))$$

$$f \mapsto \hat{f}$$

$$g \circ \eta_\Sigma \leftarrow g$$

(where $\eta_\Sigma : \Sigma \rightarrow F \Sigma = \text{List } \Sigma$ is $a \mapsto [a]$)

The bijection is “natural in Σ and (M, \cdot, e) ” (to be explained)

Binary product in a category \mathbf{C}

$(Z, Z) \rightarrow (X, Y)$ morphisms in $\mathbf{C} \times \mathbf{C}$

$Z \rightarrow X \times Y$ morphisms in \mathbf{C}

bijection

$(\mathbf{C} \times \mathbf{C})((Z, Z), (X, Y)) \cong \mathbf{C}(Z, X \times Y)$

$(f, g) \mapsto \langle f, g \rangle$

$(\pi_1 \circ h, \pi_2 \circ h) \leftarrow h$

This bijection is “natural in X, Y, Z ” (to be explained)

Exponentials in a category \mathbf{C} with binary products

$$\frac{Z \times X \rightarrow Y \text{ morphisms in } \mathbf{C}}{Z \rightarrow Y^X \text{ morphisms in } \mathbf{C}}$$

bijection

$$\mathbf{C}(Z \times X, Y) \cong \mathbf{C}(Z, Y^X)$$

$$f \mapsto \text{cur } f$$

$$\text{app} \circ (g \times \text{id}_X) \leftarrow g$$

The bijection is “natural in X, Y, Z ” (to be explained)

Adjunction

Definition. An **adjunction** between two categories **C** and **D** is specified by:

- ▶ functors $\mathbf{C} \begin{array}{c} \xrightarrow{F} \\ \xleftarrow{G} \end{array} \mathbf{D}$
- ▶ for each $X \in \mathbf{C}$ and $Y \in \mathbf{D}$ a bijection $\theta_{X,Y} : \mathbf{D}(FX, Y) \cong \mathbf{C}(X, GY)$ which is **natural in X and Y**.

$$\text{for all } \begin{cases} u : X' \rightarrow X \text{ in } \mathbf{C} \\ v : Y \rightarrow Y' \text{ in } \mathbf{D} \end{cases} \quad \text{and all } g : FX \rightarrow Y \text{ in } \mathbf{D}$$
$$X' \xrightarrow{u} X \xrightarrow{\theta_{X,Y}(g)} GY \xrightarrow{Gv} GY' = \theta_{X',Y'} \left(FX' \xrightarrow{Fu} FX \xrightarrow{g} Y \xrightarrow{v} Y' \right)$$

Adjunction

Definition. An **adjunction** between two categories **C** and **D** is specified by:

▶ functors $\mathbf{C} \begin{array}{c} \xrightarrow{F} \\ \xleftarrow{G} \end{array} \mathbf{D}$

▶ for each $X \in \mathbf{C}$ and $Y \in \mathbf{D}$ a bijection
 $\theta_{X,Y} : \mathbf{D}(F X, Y) \cong \mathbf{C}(X, G Y)$
which is **natural in X and Y**.

what has this to do with the concept of natural transformation between functors?

Hom functors

If \mathbf{C} is a locally small category, then we get a functor

$$\text{Hom}_{\mathbf{C}} : \mathbf{C}^{\text{op}} \times \mathbf{C} \rightarrow \text{Set}$$

with $\text{Hom}_{\mathbf{C}}(X, Y) \triangleq \mathbf{C}(X, Y)$ and

$$\text{Hom}_{\mathbf{C}} \left((X, Y) \xrightarrow{(f, g)} (X', Y') \right) \triangleq \mathbf{C}(X, Y) \xrightarrow{\text{Hom}_{\mathbf{C}}(f, g)} \mathbf{C}(X', Y')$$
$$\text{Hom}_{\mathbf{C}}(f, g) h \triangleq g \circ h \circ f$$

Hom functors

If \mathbf{C} is a locally small category, then we get a functor

$$\text{Hom}_{\mathbf{C}} : \mathbf{C}^{\text{op}} \times \mathbf{C} \rightarrow \text{Set}$$

with $\text{Hom}_{\mathbf{C}}(X, Y) \triangleq \mathbf{C}(X, Y)$ and

$$\text{Hom}_{\mathbf{C}} \left((X, Y) \xrightarrow{(f, g)} (X', Y') \right) \triangleq \mathbf{C}(X, Y) \xrightarrow{\text{Hom}_{\mathbf{C}}(f, g)} \mathbf{C}(X', Y')$$

$\text{Hom}_{\mathbf{C}}(f, g) h \triangleq g \circ h \circ f$

If $(f, g) : (X, Y) \rightarrow (X', Y')$ in $\mathbf{C}^{\text{op}} \times \mathbf{C}$ and $h : X \rightarrow Y$ in \mathbf{C} , then in \mathbf{C} we have $f : X' \rightarrow X$, $g : Y \rightarrow Y'$ and so $g \circ h \circ f : X' \rightarrow Y'$

Natural isomorphisms

Given functors $F, G : \mathbf{C} \rightarrow \mathbf{D}$, a **natural isomorphism** $\theta : F \cong G$ is simply an isomorphism between F and G in the functor category $\mathbf{D}^{\mathbf{C}}$.

Natural isomorphisms

Given functors $F, G : \mathbf{C} \rightarrow \mathbf{D}$, a **natural isomorphism** $\theta : F \cong G$ is simply an isomorphism between F and G in the functor category $\mathbf{D}^{\mathbf{C}}$.

Lemma. If $\theta : F \rightarrow G$ is a natural transformation and for each $X \in \mathbf{C}$, $\theta_X : FX \rightarrow GX$ is an isomorphism in \mathbf{D} , then the family of morphisms $(\theta_X^{-1} : GX \rightarrow FX \mid X \in \mathbf{C})$ gives a natural transformation $\theta^{-1} : G \rightarrow F$ which is inverse to θ in $\mathbf{D}^{\mathbf{C}}$ and hence θ is a natural isomorphism. \square

An adjunction between locally small categories \mathbf{C} and \mathbf{D} is simply a triple (F, G, θ) where

▶ $\mathbf{C} \begin{array}{c} \xrightarrow{F} \\ \xleftarrow{G} \end{array} \mathbf{D}$

▶ θ is a natural isomorphism between the functors

$$\begin{array}{ccccc}
 & & \mathbf{D}^{\text{op}} \times \mathbf{D} & & \\
 & \nearrow^{F^{\text{op}} \times \text{id}_{\mathbf{D}}} & & \searrow^{\text{Hom}_{\mathbf{D}}} & \\
 \mathbf{C}^{\text{op}} \times \mathbf{D} & & & & \mathbf{Set} \\
 & \searrow_{\text{id}_{\mathbf{C}^{\text{op}}} \times G} & & \nearrow_{\text{Hom}_{\mathbf{C}}} & \\
 & & \mathbf{C}^{\text{op}} \times \mathbf{C} & &
 \end{array}$$

and

Terminology:

Given $C \begin{array}{c} \xrightarrow{F} \\ \xleftarrow{G} \end{array} D$

is there is some natural isomorphism

$$\theta : \text{Hom}_D \circ (F^{\text{op}} \times \text{id}_D) \cong \text{Hom}_C \circ (\text{id}_{C^{\text{op}}} \times G)$$

one says

F is a **left adjoint** for G

G is a **right adjoint** for F

and writes

$$F \dashv G$$

Notation associated with an adjunction (F, G, θ)

$$\text{Given } \begin{cases} g : F X \rightarrow Y \\ f : X \rightarrow G Y \end{cases}$$

$$\text{we write } \begin{cases} \bar{g} \triangleq \theta_{X,Y}(g) : X \rightarrow G Y \\ \bar{f} \triangleq \theta_{X,Y}^{-1}(f) : F X \rightarrow Y \end{cases}$$

Thus $\bar{\bar{g}} = g$, $\bar{\bar{f}} = f$ and naturality of $\theta_{X,Y}$ in X and Y means that

$$\overline{v \circ g \circ F u} = G v \circ \bar{g} \circ u$$

Notation associated with an adjunction (F, G, θ)

The existence of θ is sometimes indicated by writing

$$\frac{FX \xrightarrow{g} Y}{X \xrightarrow{\bar{g}} GY}$$

Using this notation, one can split the naturality condition for θ into two:

$$\frac{FX' \xrightarrow{Fu} FX \xrightarrow{g} Y}{X' \xrightarrow{u} X \xrightarrow{\bar{g}} GY} \qquad \frac{FX \xrightarrow{g} Y \xrightarrow{v} Y'}{X \xrightarrow{\bar{g}} GY \xrightarrow{Gv} GY'}$$

Theorem. A category \mathbf{C} has binary products iff the diagonal functor $\Delta = \langle \text{id}_{\mathbf{C}}, \text{id}_{\mathbf{C}} \rangle : \mathbf{C} \rightarrow \mathbf{C} \times \mathbf{C}$ has a right adjoint.

Theorem. A category \mathbf{C} with binary products also has all exponentials of pairs of objects iff for all $X \in \mathbf{C}$, the functor $(-) \times X : \mathbf{C} \rightarrow \mathbf{C}$ has a right adjoint.

we'll see next time a theorem characterising adjoint functors of which the above are special cases