

Number 333



UNIVERSITY OF  
CAMBRIDGE

Computer Laboratory

## What is a categorical model of intuitionistic linear logic?

G.M. Bierman

April 1994

15 JJ Thomson Avenue  
Cambridge CB3 0FD  
United Kingdom  
phone +44 1223 763500  
<https://www.cl.cam.ac.uk/>

© 1994 G.M. Bierman

Technical reports published by the University of Cambridge  
Computer Laboratory are freely available via the Internet:

*<https://www.cl.cam.ac.uk/techreports/>*

ISSN 1476-2986

# What is a Categorical Model of Intuitionistic Linear Logic?

G.M. Bierman  
University of Cambridge Computer Laboratory

March 31, 1994

## Abstract

This paper re-addresses the old problem of providing a categorical model for Intuitionistic Linear Logic (**ILL**). In particular we compare the now standard model proposed by Seely to the lesser known one proposed by Benton, Bierman, Hyland and de Paiva. Surprisingly we find that Seely's model is *unsound* in that it does not preserve equality of proofs—we shall give some examples of equal proofs which do not seem to be modelled as equal morphisms in the category. We shall propose how to adapt Seely's definition so as to correct these problems and consider how this compares with the model due to Benton *et al.*

# 1 Intuitionistic Linear Logic

For the first part we shall consider only the multiplicative, exponential fragment of Intuitionistic Linear Logic (**MELL**). Rather than give a detailed description of the logic and associated term calculus we assume that the reader is familiar with other work [18, 5]. The sequent calculus formulation is originally due to Girard and Lafont [9] and is given below.

$$\begin{array}{c}
 \frac{}{A \vdash A} \textit{Identity} \\
 \\
 \frac{\Gamma \vdash B \quad B, \Delta \vdash C}{\Gamma, \Delta \vdash C} \textit{Cut} \\
 \\
 \frac{\Gamma \vdash A}{\Gamma, I \vdash A} (I_{\mathcal{L}}) \qquad \frac{}{\vdash I} (I_{\mathcal{R}}) \\
 \\
 \frac{\Gamma, A, B \vdash C}{\Gamma, A \otimes B \vdash C} (\otimes_{\mathcal{L}}) \qquad \frac{\Gamma \vdash A \quad \Delta \vdash B}{\Gamma, \Delta \vdash A \otimes B} (\otimes_{\mathcal{R}}) \\
 \\
 \frac{\Gamma \vdash A \quad \Delta, B \vdash C}{\Gamma, \Delta, A \multimap B \vdash C} (\multimap_{\mathcal{L}}) \qquad \frac{\Gamma, A \vdash B}{\Gamma \vdash A \multimap B} (\multimap_{\mathcal{R}}) \\
 \\
 \frac{\Gamma \vdash B}{\Gamma, !A \vdash B} \textit{Weakening} \qquad \frac{\Gamma, !A, !A \vdash B}{\Gamma, !A \vdash B} \textit{Contraction} \\
 \\
 \frac{\Gamma, A \vdash B}{\Gamma, !A \vdash B} \textit{Dereliction} \qquad \frac{! \Gamma \vdash A}{! \Gamma \vdash !A} \textit{Promotion}
 \end{array}$$

Sequents are written as  $\Gamma \vdash A$ , where  $A, B$  represent formulae and  $\Gamma, \Delta$  represent multisets of formulae. Where  $\Gamma$  represents the multiset  $A_1, \dots, A_n$ , then  $! \Gamma$  is taken to represent the multiset  $!A_1, \dots, !A_n$ .

The natural deduction presentation proved harder to formalize and early proposals [1, 15] failed to have the vital property of *closure under substitution*. A natural deduction system which has this property was given by Benton *et al.* [5] and is given below.

$$\begin{array}{c}
 [A^x] \\
 \vdots \\
 B \\
 \hline
 A \multimap B \quad (\multimap_{\mathcal{L}})_x
 \end{array}
 \qquad
 \begin{array}{c}
 \vdots \qquad \vdots \\
 A \multimap B \qquad A \\
 \hline
 B \quad (\multimap_{\mathcal{R}})
 \end{array}$$
  

$$\begin{array}{c}
 \frac{}{I} (I_{\mathcal{L}})
 \end{array}
 \qquad
 \begin{array}{c}
 \vdots \qquad \vdots \\
 A \qquad I \\
 \hline
 A \quad (I_{\mathcal{R}})
 \end{array}$$

$$\begin{array}{c}
\begin{array}{c} \vdots \\ A \end{array} \quad \begin{array}{c} \vdots \\ B \end{array} \\
\hline
A \otimes B \quad (\otimes_I)
\end{array}
\qquad
\begin{array}{c}
\begin{array}{c} [A^x] [B^y] \\ \vdots \\ A \otimes B \end{array} \quad \begin{array}{c} \vdots \\ C \end{array} \\
\hline
C \quad (\otimes_\varepsilon)_{x,y}
\end{array}$$

$$\begin{array}{c}
\begin{array}{c} \vdots \\ !B \end{array} \quad \begin{array}{c} \vdots \\ C \end{array} \\
\hline
C \quad \textit{Weakening}
\end{array}
\qquad
\begin{array}{c}
\begin{array}{c} [!B^x] [!B^y] \\ \vdots \\ !B \end{array} \quad \begin{array}{c} \vdots \\ C \end{array} \\
\hline
C \quad \textit{Contraction}_{x,y}
\end{array}$$

$$\begin{array}{c}
\begin{array}{c} \vdots \\ !B \end{array} \\
\hline
B \quad \textit{Dereliction}
\end{array}
\qquad
\begin{array}{c}
\begin{array}{c} \vdots \\ !A_1 \quad \dots \quad !A_n \end{array} \quad \begin{array}{c} \vdots \\ B \end{array} \\
\hline
!B \quad \textit{Promotion}_{x_1, \dots, x_n}
\end{array}$$

The main difference between this and earlier presentations is in the *Promotion* rule where here substitutions are ‘built-in’.

The Curry-Howard correspondence [10] provides a systematic process for attaching names, or *terms*, to proof trees from the natural deduction formulation of a given constructive logic (a clear description is given by Gallier [7]). We can apply it to get the following term assignment system for **MELL**, which rather than presenting in a tree-like fashion, we choose to present in a sequent style.

$$\begin{array}{c}
\frac{}{x: A \triangleright x: A} \textit{Identity}
\end{array}$$

$$\begin{array}{c}
\frac{\Gamma, x: A \triangleright M: B}{\Gamma \triangleright \lambda x: A. M: A \multimap B} (-\otimes_I) \qquad \frac{\Gamma \triangleright M: A \multimap B \quad \Delta \triangleright N: A}{\Gamma, \Delta \triangleright MN: B} (-\otimes_\varepsilon)
\end{array}$$

$$\begin{array}{c}
\frac{}{\triangleright *: I} (I_I) \qquad \frac{\Gamma \triangleright M: A \quad \Delta \triangleright N: I}{\Gamma, \Delta \triangleright \text{let } N \text{ be } * \text{ in } N: A} (I_\varepsilon)
\end{array}$$

$$\begin{array}{c}
\frac{\Gamma \triangleright M: A \quad \Delta \triangleright N: B}{\Gamma, \Delta \triangleright M \otimes N: A \otimes B} (\otimes_I) \qquad \frac{\Delta \triangleright M: A \otimes B \quad \Gamma, x: A, y: B \triangleright N: C}{\Gamma, \Delta \triangleright \text{let } M \text{ be } x \otimes y \text{ in } N: C} (\otimes_\varepsilon)
\end{array}$$

$$\frac{\Gamma_1 \triangleright M_1: !A_1 \quad \dots \quad \Gamma_n \triangleright M_n: !A_n \quad x_1: !A_1, \dots, x_n: !A_n \triangleright N: B}{\Gamma_1, \dots, \Gamma_n \triangleright \text{promote } M_1, \dots, M_n \text{ for } x_1, \dots, x_n \text{ in } N: !B} \textit{Promotion}$$

$$\frac{\Gamma \triangleright M: !A \quad \Delta \triangleright N: B}{\Gamma, \Delta \triangleright \text{discard } M \text{ in } N: B} \textit{Weakening}$$

$$\frac{\Delta \triangleright M: !A \quad \Gamma, x: !A, y: !A \triangleright N: B}{\Gamma, \Delta \triangleright \text{copy } M \text{ as } x, y \text{ in } N: B} \textit{Contraction}$$

$$\frac{\Gamma \triangleright M: !A}{\Gamma \triangleright \text{derelict}(M): A} \textit{Dereliction}$$

Subsequently, others have suggested different natural deduction formulations which have the property of more compact syntax for the *Promotion* rule. The reader is referred to

work by Troelstra [16] and Wadler [17] for details of two proposals.

Normalization is the process of removing ‘detours’ from a proof in natural deduction. At the level of terms it can be seen as providing a set of reduction rules, which are known as  $\beta$ -rules. For **MELL** there are six  $\beta$ -rules which are given below.

1.  $(\lambda x: A. M) N \rightsquigarrow_{\beta} M[x := N]$
2.  $\text{let } * \text{ be } * \text{ in } M \rightsquigarrow_{\beta} M$
3.  $\text{let } M \otimes N \text{ be } x \otimes y \text{ in } P \rightsquigarrow_{\beta} P[x := M, y := N]$
4.  $\text{derelict}(\text{promote } \vec{M} \text{ for } \vec{x} \text{ in } N) \rightsquigarrow_{\beta} N[\vec{x} := \vec{M}]$
5.  $\text{discard}(\text{promote } \vec{M} \text{ for } \vec{x} \text{ in } N) \text{ in } P \rightsquigarrow_{\beta} \text{discard } \vec{M} \text{ in } P$
6.  $\text{copy}(\text{promote } \vec{M} \text{ for } \vec{x} \text{ in } N) \text{ as } y, z \text{ in } P \rightsquigarrow_{\beta} \text{copy } \vec{M} \text{ as } \vec{u}, \vec{v} \text{ in}$   
 $P [ \quad y := \text{promote } \vec{u} \text{ for } \vec{x} \text{ in } N,$   
 $\quad z := \text{promote } \vec{v} \text{ for } \vec{x} \text{ in } N]$

In addition there are other term equalities: *commuting conversions*, which arise from consideration of the subformula property, as well as the those suggested by the process of cut elimination for the sequent calculus formulation.<sup>1</sup> For the purposes of this paper these need not be considered here. The interested reader is again referred to other work [6, 5].

## 2 Two Categorical Models

The fundamental idea of a categorical treatment of proof theory is that propositions should be interpreted as the objects of the category and proofs should be interpreted as morphisms. The proof rules correspond to natural transformations between appropriate hom-functors. As mentioned above, the proof theoretic setting will reveal a number of reduction rules, which can be viewed as equalities between proofs. In particular, these equalities should hold in the categorical model.

Let us fix some notation. The interpretation of a proof is represented using semantic braces,  $\llbracket - \rrbracket$ , making the usual simplification of using the same letter to represent a proposition as its interpretation. Given a term  $\Gamma \triangleright M: A$  where  $M \rightsquigarrow_{\beta} N$ , we shall write  $\Gamma \triangleright M = N: A$ .

**Definition 1** *A category,  $\mathbb{C}$ , is said to be a categorical model of a given logic,  $\mathcal{L}$ , iff*

1. *For all proofs  $\Gamma \triangleright_{\mathcal{L}} M: A$ , there is a morphism  $\llbracket M \rrbracket: \Gamma \rightarrow A$  in  $\mathbb{C}$ .*
2. *For all equalities  $\Gamma \triangleright_{\mathcal{L}} M = N: A$  it is the case that  $\llbracket M \rrbracket =_{\mathbb{C}} \llbracket N \rrbracket$  (where  $=_{\mathbb{C}}$  represents equality of morphisms in the category  $\mathbb{C}$ ).*

Given this definition we shall now consider two proposals for a categorical model of Linear Logic. Firstly that proposed by Seely [14] and secondly that of Benton *et al.* [5]. First let us recall Seely’s definition (where for clarity we have named the natural isomorphisms relating the tensor and categorical products).

**Definition 2 (Seely)** *A Seely category,  $\mathbb{C}$ , consists of:*

1. *A symmetric monoidal closed category (SMCC) with finite products, together with a comonad  $(!, \varepsilon, \delta)$ .*

---

<sup>1</sup>In fact there are other term equalities due to the interaction between our formulation of the *Promotion* rule and the fact that we are suppressing the *Exchange* rule.

2. For each object  $A$  of  $\mathbb{C}$ ,  $(!A, d_A, e_A)$  is a comonoid with respect to the tensor product.
3. There exists natural isomorphisms  $n: !A \otimes !B \xrightarrow{\sim} !(A \times B)$  and  $p: I \xrightarrow{\sim} !1$ .
4. The functor  $!$  takes the comonoid structure of the cartesian product to the comonoid structure of the tensor product.

It is instructive to consider this definition in more detail. The naturality of  $n$  amounts to the following diagram commuting for morphisms  $f: A \rightarrow C$  and  $g: B \rightarrow D$ .

$$\begin{array}{ccc}
 !A \otimes !B & \xrightarrow{n} & !(A \times B) \\
 \downarrow !f \otimes !g & & \downarrow !(f \times g) \\
 !C \otimes !D & \xrightarrow{n} & !(C \times D)
 \end{array}$$

Condition 4 (which seems to have been overlooked<sup>2</sup> by Barr [3] and Troelstra [15]) amounts to requiring that the following two diagrams commute.

$$\begin{array}{ccc}
 !A & \xrightarrow{d_A} & !A \otimes !A \\
 \searrow !\Delta & & \downarrow n \\
 & & !(A \times A)
 \end{array}
 \qquad
 \begin{array}{ccc}
 !A & \xrightarrow{e_A} & I \\
 \searrow !\top & & \downarrow p \\
 & & !1
 \end{array}$$

Now let us consider the model proposed by Benton *et al.* (the version given here is taken from my thesis [6] and is a slight adaptation from the original definition [5]).

**Definition 3** A Linear category,  $\mathbb{C}$ , consists of:

1. A SMCC,  $\mathbb{C}$ , together with:
2. A symmetric monoidal comonad  $(!, \varepsilon, \delta, m_{A,B}, m_I)$  such that
  - (a) For every free  $!$ -coalgebra  $(!A, \delta_A)$  there are two distinguished monoidal natural transformations with components  $e_A: !A \rightarrow I$  and  $d_A: !A \rightarrow !A \otimes !A^3$  which form a commutative comonoid and are coalgebra morphisms.
  - (b) Whenever  $f: (!A, \delta_A) \rightarrow (!B, \delta_B)$  is a coalgebra morphism between free coalgebras, then it is also a comonoid morphism.

Let us consider in detail the conditions in this definition. Firstly requiring that  $(!, m_{A,B}, m_I)$  is a symmetric monoidal functor amounts to the following diagrams commuting.

<sup>2</sup>Asperti and Longo [2, Lemma 5.5.4] (falsely) claim it holds automatically.

<sup>3</sup>This necessitates showing that  $!\otimes!$  and  $I$  are monoidal functors, but this is trivial and omitted.

$$\begin{array}{ccc}
!I \otimes !A & \xrightarrow{m_{I,A}} & !(I \otimes A) \\
\uparrow m_I \otimes \text{id}_{!A} & & \downarrow !(\lambda_A) \\
I \otimes !A & \xrightarrow{\lambda_{!A}} & !A \\
\end{array}
\qquad
\begin{array}{ccc}
!A \otimes !I & \xrightarrow{m_{A,I}} & !(A \otimes I) \\
\uparrow \text{id}_{!A} \otimes m_I & & \downarrow !(\rho_A) \\
!A \otimes I & \xrightarrow{\rho_{!A}} & !A \\
\end{array}$$

$$\begin{array}{ccccc}
(!A \otimes !B) \otimes !C & \xrightarrow{m_{A,B} \otimes \text{id}_{!C}} & !(A \otimes B) \otimes !C & \xrightarrow{m_{A \otimes B, C}} & !((A \otimes B) \otimes C) \\
\uparrow \alpha_{!A, !B, !C} & & & & \uparrow !(\alpha_{A, B, C}) \\
!A \otimes (!B \otimes !C) & \xrightarrow{\text{id}_{!A} \otimes m_{B,C}} & !A \otimes (B \otimes C) & \xrightarrow{m_{A, B \otimes C}} & !(A \otimes (B \otimes C)) \\
\end{array}$$

$$\begin{array}{ccc}
!A \otimes !B & \xrightarrow{m_{A,B}} & !(A \otimes B) \\
\downarrow \gamma_{A,B} & & \downarrow !(\gamma_{A,B}) \\
!B \otimes !A & \xrightarrow{m_{B,A}} & !(B \otimes A) \\
\end{array}$$

Requiring that  $\varepsilon$  is a monoidal natural transformation amounts to the following two commuting diagrams.



$$\begin{array}{ccc}
!A \otimes !B & \xrightarrow{m_{A,B}} & !(A \otimes B) \\
\searrow \varepsilon_{A \otimes B} & & \downarrow \varepsilon_{A \otimes B} \\
& & A \otimes B
\end{array}
\qquad
\begin{array}{ccc}
I & & \\
\downarrow m_I & \cong & \\
!I & \xrightarrow{\varepsilon_I} & I
\end{array}$$

Requiring that  $\delta$  is a monoidal natural transformation amounts to the following two commuting diagrams.


$$\begin{array}{ccc}
!A \otimes !B & \xrightarrow{m_{A,B}} & !(A \otimes B) \\
\downarrow \delta_A \otimes \delta_B & & \downarrow \delta_{A \otimes B} \\
!!A \otimes !!B & \xrightarrow{m_{!A, !B}} & !(A \otimes B) \\
& & \downarrow !m_{A,B} \\
& & !!(A \otimes B)
\end{array}$$

$$\begin{array}{ccc}
I & \xrightarrow{m_I} & !I \\
\downarrow m_I & & \downarrow \delta_I \\
!I & \xrightarrow{!m_I} & !!I
\end{array}$$


Requiring that  $e_A: !A \rightarrow I$  is a monoidal natural transformation amounts to requiring that the following three diagrams commute, for any morphism  $f: A \rightarrow B$ .




Requiring that  $d_A: !A \rightarrow !A \otimes !A$  is a monoidal natural transformation amounts to requiring that the following three diagrams commute, for all  $f: A \rightarrow B$ .




Requiring that  $(!A, d_A, e_A)$  forms a commutative comonoid amounts to requiring that the following three diagrams commute.







Requiring that  $e_A$  is a coalgebra morphism amounts to requiring that the following diagram commutes.



Requiring that  $d_A$  is a coalgebra morphism amounts to requiring that the following diagram commutes.



Finally all coalgebra morphisms between *free* coalgebras are also comonoid morphisms. Thus given a coalgebra morphism  $f$ , between the free coalgebras  $(!A, \delta_A)$  and  $(!B, \delta_B)$ , i.e. which makes the following diagram commute.



Then it is also a comonoid morphism between the comonoids  $(!A, e_A, d_A)$  and  $(!B, e_B, d_B)$ , i.e. it makes the following diagram commute.

$$\begin{array}{ccc}
& !A & \xrightarrow{d_A} & !A \otimes !A \\
& \swarrow e_A & & \downarrow f \otimes f \\
I & & f & \\
& \searrow e_B & & \downarrow \\
& !B & \xrightarrow{d_B} & !B \otimes !B
\end{array}$$

These amount to some strong conditions on the model and some of their consequences are explored in my thesis. It is, however, reasonably straightforward to show the following.

**Theorem 1** *A Linear category,  $\mathbb{C}$ , is a categorical model for **MELL**.*

**Proof.** The first condition is proved by a trivial induction on the structure of the proof  $\Gamma \triangleright M : A$ . The second condition is proved by checking that the six  $\beta$ -rules from earlier. ■

The main difference between these two models is that a Seely category critically needs categorical products to model the exponential (!). Consider the interpretation of the *Promotion* rule. With a Seely category this is interpreted:

$$\begin{aligned}
& \llbracket \Gamma_1, \dots, \Gamma_n \triangleright \text{promote } M_1, \dots, M_n \text{ for } x_1, \dots, x_n \text{ in } N : !B \rrbracket \\
& \stackrel{\text{def}}{=} \llbracket \Gamma_1 \triangleright M_1 : !A_1 \rrbracket \otimes \dots \otimes \llbracket \Gamma_n \triangleright M_n : !A_n \rrbracket ; n ; \delta ; !n^{-1} ; !(\llbracket x_1 : !A_1, \dots, x_n : !A_n \triangleright N : B \rrbracket)
\end{aligned}$$

With a Linear category this is interpreted:

$$\begin{aligned}
& \llbracket \Gamma_1, \dots, \Gamma_n \triangleright \text{promote } M_1, \dots, M_n \text{ for } x_1, \dots, x_n \text{ in } N : !B \rrbracket \\
& \stackrel{\text{def}}{=} \llbracket \Gamma_1 \triangleright M_1 : !A_1 \rrbracket \otimes \dots \otimes \llbracket \Gamma_n \triangleright M_n : !A_n \rrbracket ; \delta \otimes \dots \otimes \delta ; m ; !(\llbracket x_1 : !A_1, \dots, x_n : !A_n \triangleright N : B \rrbracket)
\end{aligned}$$

Let us consider whether a Seely category is a categorical model for **MELL**. Seely showed that the first requirement is satisfied.

**Proposition 1 (Seely)** *Given a Seely category,  $\mathbb{C}$ , for all proofs  $\Gamma \triangleright M : A$  there is a morphism  $\llbracket M \rrbracket : \Gamma \rightarrow A$  in  $\mathbb{C}$ .*

However the second condition is not satisfied.

**Fact 1** *Given a Seely category,  $\mathbb{C}$ , it is not the case that for all term equalities  $\Gamma \triangleright M = N : A$  that  $\llbracket M \rrbracket =_{\mathbb{C}} \llbracket N \rrbracket$ .<sup>4</sup>*

A counter-example is the sixth  $\beta$ -rule from earlier. In fact we only need use a simplified version where the promoted term,  $N$ , has only one free variable.

<sup>4</sup>It should be noted that the term equalities were not generally known when Seely proposed his model.

$\Gamma \triangleright$  copy (promote  $M$  for  $x$  in  $N$ ) as  $y, z$  in  $P$   
 $=$  copy  $M$  as  $x', x''$  in  $P[y := \text{promote } x' \text{ for } x \text{ in } N, z := \text{promote } x'' \text{ for } x \text{ in } N]: C$

This term equality implies the *same commuting diagram* for a Linear category as for a Seely category.

$$\begin{array}{ccccccc}
\Gamma & \xrightarrow{m} & !A & \xrightarrow{\delta} & !!A & \xrightarrow{!n} & !B \\
& & \downarrow d & & & & \downarrow d \\
& & !A \otimes !A & \xrightarrow{\delta \otimes \delta} & !!A \otimes !!A & \xrightarrow{!n \otimes !n} & !B \otimes !B \xrightarrow{p} C
\end{array} \tag{1}$$

For a Linear category we can complete the diagram in the following way.

$$\begin{array}{ccccccc}
\Gamma & \xrightarrow{m} & !A & \xrightarrow{\delta} & !!A & \xrightarrow{!n} & !B \\
& & \downarrow d & & \downarrow d & & \downarrow d \\
& & !A \otimes !A & \xrightarrow{\delta \otimes \delta} & !!A \otimes !!A & \xrightarrow{!n \otimes !n} & !B \otimes !B \xrightarrow{p} C
\end{array}$$

The left hand square commutes by the condition that all free coalgebra morphisms are comonoid morphisms. The right hand square commutes by naturality of  $d$ . Unfortunately it is not clear how to make diagram 1 commute for a Seely category.

At this stage we might try adding the condition that all (free) coalgebra morphisms are comonoid morphisms to Seely's definition. This proves still to be incomplete as we find that neither the sixth nor the fifth term equalities are modelled correctly in the cases when the promoted term,  $N$ , has zero or more than one free variable. One might be further tempted to add additional *ad-hoc* conditions to make a Seely category a model for **MELL**. However, as shown in my thesis, this is by no means simple and rather it would seem more prudent to consider a more abstract view. Let us consider some of the motivation behind the Seely construction.

First we shall recall a construction, the dual of which (i.e. that generated by a monad) is known as the "Kleisli category" [13, Page 143].

**Definition 4** Given a comonad  $(!, \varepsilon, \delta)$  on a category  $\mathbb{C}$ , we take all the objects  $A$  in  $\mathbb{C}$  and for each morphism  $f: !A \rightarrow B$  in  $\mathbb{C}$  we take a new morphism  $\hat{f}: A \rightarrow B$ . The objects and morphisms form the co-Kleisli category  $\mathbb{C}_!$ , where the composition of the morphisms  $\hat{f}: A \rightarrow B$  and  $\hat{g}: B \rightarrow C$  is defined by the following:

$$\hat{f}; \hat{g} \stackrel{\text{def}}{=} (\delta_A; \widehat{!f}; g)$$

Our interest in this construction is that this has strong similarities with the Girard translation [8] of Intuitionistic Logic into Intuitionistic Linear Logic where the intuitionistic implication is decomposed:  $(A \supset B)^\circ \stackrel{\text{def}}{=} !(A^\circ) \multimap B^\circ$ . In fact, as first shown by Seely [14], the co-Kleisli construction can be thought of as a categorical equivalent of the Girard translation in the following sense.

**Proposition 2 (Seely)** *Given a Seely category,  $\mathbb{C}$ , the co-Kleisli category  $\mathbb{C}_!$  is cartesian closed.*

**Proof.** (Sketch) Given two objects  $A$  and  $B$  their exponent is defined to be  $!A \multimap B$ . Then we have the following sequence of isomorphisms.

$$\begin{aligned}
\mathbb{C}_!(A \times B, C) &\cong \mathbb{C}(!A \times B, C) && \text{By definition} \\
&\cong \mathbb{C}(!A \otimes !B, C) && \text{By use of the } n \text{ isomorphism} \\
&\cong \mathbb{C}(!A, !B \multimap C) && \text{By } \mathbb{C} \text{ having a closed structure} \\
&\cong \mathbb{C}_!(A, !B \multimap C) && \text{By definition}
\end{aligned}$$

■

We know from Kleisli's construction that we have the following adjunction, where  $G$  is the functor defined by  $g: A \rightarrow B \mapsto (\varepsilon; \widehat{g})$  and  $F$  is the functor defined by  $\hat{f}: A \rightarrow B \mapsto \delta_A; !f$ .

$$\begin{array}{ccc}
& \mathbb{C}_! & \\
F \downarrow & \dashv & \uparrow G \\
& \mathbb{C} &
\end{array}$$

Seely's model arises from the desire to make the co-Kleisli category a cartesian closed category (CCC), which is achieved by including the  $n$  and  $p$  natural isomorphisms. This means that there is an adjunction between a SMCC ( $\mathbb{C}$ ) and a CCC ( $\mathbb{C}_!$ ). As a CCC is trivially a SMCC, there is then an adjunction between two SMCCs. We might expect that this is a *monoidal adjunction*.

**Definition 5** *An adjunction  $\langle F, G, \eta, \epsilon \rangle: \mathbb{C} \rightarrow \mathbb{D}$  is said to be a monoidal adjunction when  $F$  and  $G$  are monoidal functors and  $\eta$  and  $\epsilon$  are monoidal natural transformations.*

Let us now state a new definition for a Seely-style category and then investigate some of its properties.

**Definition 6** *A new-Seely category,  $\mathbb{C}$ , consists of*

1. a SMCC,  $\mathbb{C}$ , with finite products, together with
2. a comonad,  $(!, \varepsilon, \delta)$ , and
3. two natural isomorphisms,  $n: !A \otimes !B \xrightarrow{\sim} !(A \times B)$  and  $p: I \xrightarrow{\sim} !1$

*such that the adjunction,  $\langle F, G, \eta, \varepsilon \rangle$ , between  $\mathbb{C}$  and  $\mathbb{C}_!$  is a monoidal adjunction.*

Assuming that  $F$  is monoidal gives us the following morphism and natural transformation:

$$\begin{aligned}
m_I: I &\rightarrow F1 \\
m_{A,B}: FA \otimes FB &\rightarrow F(A \times B)
\end{aligned}$$

Assuming that  $G$  is monoidal gives us the following morphism and natural transformation;

$$\begin{aligned}
m'_1: 1 &\rightarrow GI \\
m'_{A,B}: GA \times GB &\rightarrow G(A \otimes B)
\end{aligned}$$

By assumption  $\varepsilon$  and  $\eta$  are monoidal natural transformations.

It is easy to see that  $m_I$  is Seely's morphism  $p$  and  $m_{A,B}$  is Seely's natural transformation  $n$ . In fact, we can define their inverses:

$$\begin{aligned} m_I^{-1} &\stackrel{\text{def}}{=} Fm'_1; \varepsilon_I: F1 \rightarrow I \\ m_{A,B}^{-1} &\stackrel{\text{def}}{=} F(\eta_A \times \eta_B); Fm'_{FA,FB}; \varepsilon_{FA \otimes FB}: F(A \times B) \rightarrow FA \otimes FB \end{aligned}$$

Hence the monoidal adjunction itself provides the isomorphisms  $!A \otimes !B \cong !(A \times B)$  and  $I \cong !1$ . As the co-Kleisli category is a CCC it has a trivial commutative comonoid structure,  $(A, \Delta, \top)$ , on all objects  $A$ . We can use this and the natural transformations arising from the monoidal adjunction to define a comonoid structure,  $(F(A), d, e)$ , on the objects of  $\mathbb{C}$ :

$$\begin{aligned} d &\stackrel{\text{def}}{=} F(\Delta); m_{A,A}^{-1}: F(A) \rightarrow F(A) \otimes F(A) \\ e &\stackrel{\text{def}}{=} F(\top); m_I^{-1}: F(A) \rightarrow I \end{aligned}$$

It is easy to see that these definitions amount to condition 4 of Seely's original definition. Thus there is at least as much structure as in Seely's original definition but with the extra structure of the monoidal adjunction. Some consequences of this adjunction are given in the following lemma.

**Lemma 1** *Given a new-Seely category,  $\mathbb{C}$ , the following facts hold:*

1. *The induced comonad  $(FG, F\eta_G, \varepsilon)$  on  $\mathbb{C}$  is a monoidal comonad  $(FG, F\eta_G, \varepsilon, m_{A,B}, m_I)$ .*
2. *The comonoid morphisms  $e: FG(A) \rightarrow I$  and  $d: FG(A) \rightarrow FG(A) \otimes FG(A)$  are monoidal natural transformations.*
3. *The comonoid morphisms  $e: FG(A) \rightarrow I$  and  $d: FG(A) \rightarrow FG(A) \otimes FG(A)$  are coalgebra morphisms.*
4. *If  $f: (FG(A), F\eta_{GA}) \rightarrow (FG(B), F\eta_{GB})$  is a coalgebra morphism then it is also a comonoid morphism.*

**Proof.** For part 1 we take the following definitions:

$$\begin{aligned} m_I &\stackrel{\text{def}}{=} m_I; Fm'_1: I \rightarrow FG(I) \\ m_{A,B} &\stackrel{\text{def}}{=} m_{GA,GB}; Fm'_{A,B}: FG(A) \otimes FG(A) \rightarrow FG(A \otimes B) \end{aligned}$$

The rest of the lemma holds by construction. ■

**Corollary 1** *Every new-Seely category is a Linear category.*

(It is clear that the converse is not true, as the Linear category need not have finite products.) We can hence show that a new-Seely category is a sound model for the **MELL**.

**Theorem 2** *A new-Seely category,  $\mathbb{C}$ , is a categorical model for **MELL***

### 3 Including the additives

Now we shall consider the whole of **ILL** by adding the additive connectives to **MELL**. Logically these are given by the following sequent calculus rules (we shall ignore the additive units).

$$\begin{array}{c}
\frac{\Gamma, A \vdash C}{\Gamma, A \times B \vdash C} (\times_{\mathcal{L}-1}) \qquad \frac{\Gamma, B \vdash C}{\Gamma, A \times B \vdash C} (\times_{\mathcal{L}-2}) \\
\\
\frac{\Gamma \vdash A \quad \Gamma \vdash B}{\Gamma \vdash A \times B} (\times_{\mathcal{R}}) \\
\\
\frac{\Gamma, A \vdash C \quad \Gamma, B \vdash C}{\Gamma, A \oplus B \vdash C} (\oplus_{\mathcal{L}}) \\
\\
\frac{\Gamma \vdash A}{\Gamma \vdash A \oplus B} (\oplus_{\mathcal{R}-1}) \qquad \frac{\Gamma \vdash B}{\Gamma \vdash A \oplus B} (\oplus_{\mathcal{R}-2})
\end{array}$$

There are a number of ways of formulating the additives in a natural deduction system which are discussed in my thesis. However, this is not the place for such discussions and we shall simply take the term assignment system which is familiar from that of the  $\lambda$ -calculus. The term assignment rules as well as the  $\beta$ -rules for the additives are given below.

$$\begin{array}{c}
\frac{\Gamma \triangleright M : A \quad \Gamma \triangleright N : B}{\Gamma \triangleright \langle M, N \rangle : A \times B} (\times_{\mathcal{I}}) \\
\\
\frac{\Gamma \triangleright M : A \times B}{\Gamma \triangleright \text{fst}(M) : A} (\times_{\mathcal{E}-1}) \qquad \frac{\Gamma \triangleright M : A \times B}{\Gamma \triangleright \text{snd}(M) : B} (\times_{\mathcal{E}-2}) \\
\\
\frac{\Gamma \triangleright M : A}{\Gamma \triangleright \text{inl}(M) : A \oplus B} (\oplus_{\mathcal{I}-1}) \qquad \frac{\Gamma \triangleright M : B}{\Gamma \triangleright \text{inr}(M) : A \oplus B} (\oplus_{\mathcal{I}-2}) \\
\\
\frac{\Delta \triangleright M : A \oplus B \quad \Gamma, x : A \triangleright N : C \quad \Gamma, y : B \triangleright P : C}{\Gamma, \Delta \triangleright \text{case } M \text{ of } \text{inl}(x) \rightarrow N \parallel \text{inr}(y) \rightarrow P : C} (\oplus_{\mathcal{E}})
\end{array}$$

$$\begin{array}{c}
\text{fst}(\langle M, N \rangle) \rightsquigarrow_{\beta} M \\
\text{snd}(\langle M, N \rangle) \rightsquigarrow_{\beta} N \\
\text{case}(\text{inl}(M)) \text{ of } \text{inl}(x) \rightarrow N \parallel \text{inr}(y) \rightarrow P \rightsquigarrow_{\beta} N[x := M] \\
\text{case}(\text{inr}(M)) \text{ of } \text{inl}(x) \rightarrow N \parallel \text{inr}(y) \rightarrow P \rightsquigarrow_{\beta} P[y := M]
\end{array}$$

To model these additive connectives we shall add finite products and coproducts to a Linear category and finite coproducts to a new-Seely category. As might be expected both models are sound.

**Theorem 3** *Both a new-Seely category with finite coproducts and a Linear category with finite products and coproducts, are models for **ILL**.*

Somewhat surprisingly, we find that the so-called Seely isomorphisms (n and p) exist in a Linear category with products.

**Lemma 2** *Given a Linear category with finite products we can define the following natural isomorphisms:*

$$\begin{aligned}
n &\stackrel{\text{def}}{=} \delta \otimes \delta; m_{!A, !B}; !(\Delta); !((\text{id} \otimes e_B) \times (e_A \otimes \text{id})); !(\rho \times \lambda); !(\varepsilon \times \varepsilon); !A \otimes !B \rightarrow !(A \times B) \\
n^{-1} &\stackrel{\text{def}}{=} d_{A \times B}; !\text{fst} \otimes !\text{snd}; !(A \times B) \rightarrow !A \otimes !B \\
p &\stackrel{\text{def}}{=} m_I; !\top: I \rightarrow !1 \\
p^{-1} &\stackrel{\text{def}}{=} e_1: !1 \rightarrow I
\end{aligned}$$

Thus the co-Kleisli category associated with a Linear category is also a CCC. Given our earlier calculations we might consider the adjunction between a Linear category and its co-Kleisli category, where we find the following holds.

**Lemma 3** *The adjunction between a Linear category,  $\mathbb{C}$ , with finite products and its co-Kleisli category,  $\mathbb{C}_!$ , is a monoidal adjunction.*

Thus when considering the complete intuitionistic fragment, the new-Seely and Linear categories are equivalent.

So far we have followed others [14, 11] and only considered whether the co-Kleisli category  $\mathbb{C}_!$  generated by the comonad is cartesian closed. It should be noted that alternatively one can consider the full Eilenberg-Moore category of coalgebras ( $\mathbb{C}^!$ ) instead. In other work [5, 6], various subcategories of  $\mathbb{C}^!$  are shown to be cartesian closed. An important feature of these (sub)categories is that the underlying category  $\mathbb{C}$  need *not* necessarily have products, in contrast to the situation for  $\mathbb{C}_!$ . The interested reader is referred to these other works.

An interesting question is whether the co-Kleisli category  $\mathbb{C}_!$  has an induced coproduct structure given a strong coproduct structure in  $\mathbb{C}$ . Seely [14] showed that  $\mathbb{C}_!$  does not have a *strong* coproduct structure. It is possible however to identify a *weak* coproduct structure. We use the following well-known fact about the co-Kleisli category [12, Corollary 6.9].

**Fact 2** *The co-Kleisli category of a comonad is equivalent to the full subcategory of the category of coalgebras consisting of the free coalgebras.*

**Lemma 4** *Given two free coalgebras  $(!A, \delta_A)$  and  $(!B, \delta_B)$ , we define their coproduct to be  $(!(A \oplus B), \delta_{!A \oplus !B})$ . We define the injection morphisms to be  $\text{inl} \stackrel{\text{def}}{=} \delta_A; !\text{inl}: !A \rightarrow !(A \oplus B)$  and  $\text{inr} \stackrel{\text{def}}{=} \delta_B; !\text{inr}: !B \rightarrow !(A \oplus B)$ , which are (free) coalgebra morphisms. Given two (free) coalgebra morphisms  $f: !A \rightarrow !C$  and  $g: !B \rightarrow !C$ , then the morphism  $(![f, g]; !\varepsilon_C): !(A \oplus B) \rightarrow !C$  is a (free) coalgebra morphism and makes a coproduct diagram commute.*

**Proof.** By simple construction. ■

## 4 Conclusions

In this paper we have considered the definition of a categorical model for **ILL**. Surprisingly, Seely's now standard definition [14] was shown to be unsound, in that it does not model all equal proofs with equal morphisms. A model given in our earlier work [5] was shown

to be sound. We have also considered a method for improving Seely's original definition so as to be sound. In fact both (sound) models turn out to be equivalent.

Lafont [11] also proposed a categorical model for **ILL**, which amounts to requiring an adjunction between a SMCC and a category of commutative comonoids. In my thesis [6] it is shown that this model is a categorical model of **ILL** by proving that every Lafont category is a Linear category.

In Lemma 1 it was proved that a monoidal adjunction between a particular SMCC (a new-Seely category) and CCC (its co-Kleisli category) yielded the structure of a Linear category. Lemma 3 shows that a Linear category also has the structure of a monoidal adjunction between it (a SMCC) and its associated co-Kleisli category (a CCC). Thus the notion of a Linear category is in some senses equivalent to the existence of a monoidal adjunction between a SMCC and a CCC. This was observed by Benton [4] in his work on a mixed linear and non-linear  $\lambda$ -calculus, where the calculus is modelled by a monoidal adjunction between a SMCC and a CCC.

## Acknowledgements

This paper is taken in part from my PhD thesis. I should like to thank Valeria de Paiva, Martin Hyland and Nick Benton for their help and useful comments on both my thesis and this paper. This work was originally funded by a studentship from the Cambridge Philosophical Society. I am currently funded by research fellowships from SERC and Wolfson College, Cambridge.

## References

- [1] S. Abramsky. Computational interpretations of linear logic. *Theoretical Computer Science*, 111(1-2):3-57, 1993. Previously Available as Department of Computing, Imperial College Technical Report 90/20, 1990.
- [2] A. Asperti and G. Longo. *Categories, Types and Structures (An Introduction to Category Theory for the Working Computer Scientist)*. MIT Press, Cambridge, Mass., 1991.
- [3] M. Barr. \*-autonomous categories and linear logic. *Mathematical Structures in Computer Science*, 1:159-178, 1991.
- [4] P.N. Benton. A mixed linear and non-linear  $\lambda$ -calculus. Unpublished Manuscript, 1993.
- [5] P.N. Benton, G.M. Bierman, V.C.V. de Paiva, and J.M.E. Hyland. Term assignment for intuitionistic linear logic. Technical Report 262, Computer Laboratory, University of Cambridge, August 1992.
- [6] G.M. Bierman. *On Intuitionistic Linear Logic*. PhD thesis, Computer Laboratory, University of Cambridge, December 1993.
- [7] J. Gallier. Constructive logics part I: A tutorial on proof systems and typed  $\lambda$ -calculi. *Theoretical Computer Science*, 110(2):249-339, March 1993.
- [8] J.-Y. Girard. Linear logic. *Theoretical Computer Science*, 50:1-101, 1987.

- [9] J.-Y. Girard and Y. Lafont. Linear logic and lazy computation. In *Proceedings of TAPSOFT 87*, volume 250 of *Lecture Notes in Computer Science*, pages 52–66, 1987. Previously Available as INRIA Report 588, 1986.
- [10] W.A. Howard. The formulae-as-types notion of construction. In J.R. Hindley and J.P. Seldin, editors, *To H.B. Curry: Essays on combinatory logic, lambda calculus and formalism*. Academic Press, 1980.
- [11] Y. Lafont. The linear abstract machine. *Theoretical Computer Science*, 59:157–180, 1988. Corrections *ibid.* 62:327–328, 1988.
- [12] J. Lambek and P.J. Scott. *Introduction to higher order categorical logic*, volume 7 of *Cambridge studies in advanced mathematics*. Cambridge University Press, 1987.
- [13] S. Mac Lane. *Categories for the Working Mathematician*, volume 5 of *Graduate Texts in Mathematics*. Springer Verlag, 1971.
- [14] R.A.G. Seely. Linear logic, \*-autonomous categories and cofree algebras. In *Conference on Categories in Computer Science and Logic*, volume 92 of *AMS Contemporary Mathematics*, pages 371–382, June 1989.
- [15] A.S. Troelstra. *Lectures on Linear Logic*, volume 29 of *Lecture Notes*. CSLI, 1992.
- [16] A.S. Troelstra. Natural deduction for intuitionistic linear logic. Technical Report ML-93-09, Institute for Logic, Language and Computation, University of Amsterdam, May 1993.
- [17] P. Wadler. A syntax for linear logic. In *Mathematical Foundations of Programming Semantics*, Lecture Notes in Computer Science. Springer Verlag, 1993.
- [18] P. Wadler. A taste of linear logic. In *Symposium on the Mathematical Foundations of Computer Science*, volume 711 of *Lecture Notes in Computer Science*, pages 185–210. Springer Verlag, August 1993.